

ON A THEOREM OF FALTINGS ON FORMAL FUNCTIONS

PAOLA BONACINI, ALESSIO DEL PADRONE, AND MICHELE NESCI

ABSTRACT. In 1980, Faltings proved, by deep local algebra methods, a local result regarding formal functions which has the following global geometric fact as a consequence. *Theorem.* — *Let k be an algebraically closed field (of any characteristic). Let Y be a closed subvariety of a projective irreducible variety X defined over k . Assume that $X \subset \mathbb{P}^n$, $\dim(X) = d > 2$ and Y is the intersection of X with r hyperplanes of $\mathbb{P}^n$, with $r \leq d - 1$. Then, every formal rational function on X along Y can be (uniquely) extended to a rational function on X .* Due to its importance, the aim of this paper is to provide two elementary global geometric proofs of this theorem.

INTRODUCTION

The aim of this work is to give two elementary global geometric proofs of the following Theorem 1, which is a consequence of a local result of Faltings [Fal80] by means of the general local-global philosophy explained in [Gro05]. Faltings original proof is not so easy to follow, and it is also not immediate that what he proved implies Theorem 1, which is, on the other hand, useful for the applications. Hence we think that giving elementary arguments could be of interest.

Theorem 1. *Let k be an algebraically closed field (of any characteristic). Let Y be a closed subvariety of a projective irreducible subvariety X defined over k . Assume that $X \subset \mathbb{P}^n$, $\dim(X) = d > 2$ and Y is the intersection of X with r hyperplanes of $\mathbb{P}^n$, with $r \leq d - 1$. Then Y is G_3 in X .*

If Y is a complete intersection of X with the hyperplanes H_i (i.e. $\dim(Y) = d - r$), this result was already proved geometrically by Hirsonaka and Matsumura [HM68, (4.3), (3.5)] (see also [Băd04, 9.25]). Faltings' result, as formulated in Theorem 1, is indeed useful in some applications. For instance, Bădescu used it in an essential way to prove a relevant strengthening of Fulton-Hansen connectedness Theorem (see [Băd96, (0.1)], cf. also [Băd04, Chapter 11], or also Example 2 below).

In this work we present two global geometric proofs of 1. Both proofs use, repeatedly, as a key tool a Theorem of Hironaka and Matsumura ([HM68, (2.7)], see also Theorem 2 below) whose proof makes essential use of Grothendieck’s existence Theorem [Gro61, (5.1.4)]. Indeed this result, under the hypotheses of Theorem 1 and by means of geometric constructions yielding suitable morphisms, reduces the problem to the case of closed connected subvarieties of projective spaces of positive dimension. This case is well known by another fundamental Theorem of Hironaka and Matsumura ([HM68, (3.3)], see also Theorem 3 below). The first one is inspired by the proof of Hironaka and Matsumura of the result, for the complete intersection case, mentioned above; in fact what we do is to provide the necessary extra arguments (not completely trivial) in order to make Hironaka-Matsumura’s proof work. As far as the second proof is concerned, we use a standard construction (involving a certain incidence variety).

The first section is devoted to recall some basic facts of formal geometry and then to explain the common strategy of the two proofs, which are based on two “projective geometry style” constructions. The first proof is given in section 2, and the second is presented in section 3.

Acknowledgments. We want to thank Professor Lucian Bădescu who proposed this problem to us and supervised our work, giving also several useful comments and suggestions. This project started during our stay in the PRAGMATIC 2006 held at the Department of Mathematics of the University of Catania. We would also like to thank this institution for their warm hospitality and for the excellent working atmosphere.

1. BACKGROUND MATERIAL

The main reference is the original work of Hironaka and Matsumura [HM68], further material, together with a gentle introduction to “formal geometry”, can be found in the textbook [Băd04].

Let $\mathcal{Z}$ be a formal scheme (see [Gro60], cf. also [Har70, Har77, Băd04]). The ring of formal rational functions on $\mathcal{Z}$, $K(\mathcal{Z})$, is defined as follows: for any affine open subset U of $\mathcal{Z}$, let $[\mathcal{O}_{\mathcal{Z}}(U)]_0$ be the total ring of fractions of $\mathcal{O}_{\mathcal{Z}}(U)$, and let $\mathcal{M}_{\mathcal{Z}}$ be the sheaf associated to the presheaf on $\mathcal{Z}$ defined by $U \mapsto [\mathcal{O}_{\mathcal{Z}}(U)]_0$. Then $K(\mathcal{Z}) = H^0(\mathcal{Z}, \mathcal{M}_{\mathcal{Z}})$. If $\mathcal{Z}$ is an ordinary scheme, $K(\mathcal{Z})$ is nothing but the usual ring of rational functions of $\mathcal{Z}$.

As a special case, for any pair (X, Y) with X a locally noetherian scheme and Y a connected closed subscheme of X , we can consider

the formal completion $\mathcal{Z} := X_{/Y}$ of X along Y . Then the completion morphism $X_{/Y} \rightarrow X$ gives rise to the canonical homomorphisms $H^0(X, \mathcal{O}_X) \rightarrow H^0(X_{/Y}, \mathcal{O}_{X_{/Y}})$ and $K(X) \rightarrow K(X_{/Y})$.

We recall that for any open neighbourhood U of Y in X we have an isomorphism of formal schemes $U_{/Y} \cong X_{/Y}$. Moreover: $X_{/Y} \cong X_{/(Y)_{\text{red}}}$, that is $X_{/Y}$ depends just on the *closed subset* Y .

By [HM68, Remark p. 57], when X is a reduced algebraic scheme (for example an algebraic variety), then $K(X_{/Y})$ is a finite direct product of fields. If Y is the disjoint union of two closed subsets Y_1 and Y_2 , then $K(X_{/Y}) = K(X_{/Y_1}) \times K(X_{/Y_2})$. Hence, in order $K(X_{/Y})$ to be a field we must have Y connected. Conversely, if X is an irreducible normal projective variety, and Y is connected, then $K(X_{/Y})$ is a field.

Examples 1. As basic examples we consider the case of (connected) subvarieties Y of $X = \mathbb{P}^n$, the complex projective space.

- a) For $Y = P \in X$ a point, say over $k = \mathbb{C}$, we have $X_{/Y} = \mathbb{P}_{/P}^n \cong \mathbb{A}_{/(0, \dots, 0)}^n$, and hence $K(\mathbb{P}_{/P}^n) \cong \mathbb{C}((x_1, \dots, x_n))$ (the field of fraction of the ring of formal power series $\mathbb{C}[[x_1, \dots, x_n]]$).
- b) In case Y is connected and positive dimensional, a fundamental result of Hironaka and Matsumura, quoted below (see Theorem 3), asserts that $K(\mathbb{P}_{/Y}^n) \cong K(\mathbb{P}^n) = \mathbb{C}(x_1, \dots, x_n)$.

A basic property of the ring of formal rational functions is given by the following useful formula due to Hironaka and Matsumura (see [HM68, (2.7)], cf. also [Băd04, 9.11]):

Theorem 2. *Let $f: X' \rightarrow X$ be a proper surjective morphism of irreducible algebraic varieties, and let $Y \subseteq X$ be a closed subvariety of X , then the canonical homomorphism*

$$[K(X') \otimes_{K(X)} K(X_{/Y})]_0 \rightarrow K(X'_{/f^{-1}(Y)})$$

is an isomorphism.

Definition 1. Let X be a scheme, and let Y be a closed subscheme of X . Following [HM68, (2.9)] (see also [Har70, V] [Băd04, 9.12]), we say that: Y is *G2* in X if $K(X_{/Y})$ is a finite module over $K(X)$; Y is *G3* in X if the canonical map $K(X) \rightarrow K(X_{/Y})$ is an isomorphism.

Remark 1. We recall some elementary facts from [HM68, (2.10)] (see also [Har70, V] and [Băd04, Chapter 9]).

- a) If X is connected and complete over an algebraically closed field, then $K(X_{/Y})$ is a finite direct product of fields. Hence, if Y is *G3* in X , then Y is necessarily connected. Moreover, *G3* $\Rightarrow$ *G2*.

- b) Theorem 2 easily implies that, if Y a closed subvariety of an irreducible variety X , then $(X', f^{-1}(Y))$ is $G3$ if and only if (X, Y) is $G3$ for every proper surjective morphism $f: X' \rightarrow X$ from an irreducible variety X' . See [HM68, (2.7)], cf. also [Har70, V] and [Băd04, 9.9, 9.13(i)].
- c) The following elementary fact will be useful. Let X be an irreducible variety with two “nested” closed subsets $Y_2 \subset Y_1 \subset X$. Assuming $K(X_{/Y_1})$ a field, then (X, Y_1) is $G3$ if (X, Y_2) is $G3$. Indeed:

$$\begin{array}{ccc}
 K(X) & \xrightarrow{\cong} & K(X_{/Y_2}) \\
 & \searrow & \nearrow \\
 & K(X_{/Y_1}) &
 \end{array}$$

The following fundamental Theorem, also due to Hironaka and Matsumura (see [HM68, (3.3)]), completely explains what happens in case Y is a closed subscheme of a projective space $X = \mathbb{P}^n$ (as already remarked, in order Y to be $G3$ in X , we must have Y connected).

Theorem 3. *Let Y be a connected closed subscheme of $X = \mathbb{P}^n$. Then, Y is $G3$ in X if (and only if) $\dim(Y) > 0$.*

Remark 2. Let X be a projective irreducible variety defined over the field of complex numbers $k = \mathbb{C}$, and let Y be a connected positive dimensional closed subvariety of X . Using results of Chow and Serre’s GAGA one can show that for every connected open subset U of X (in the complex topology of X) containing Y , one has the following inclusions (see e.g. [Băd04], Chapter 10)

$$K(X) \subseteq \mathcal{M}(U) \subseteq K(X_{/Y}).$$

In particular, if Y is $G3$ in X then one gets that $K(X) = \mathcal{M}(U)$. Therefore Theorem 3 implies the following analytic result of Severi-Barth: For every closed connected subvariety Y of $\mathbb{P}_{\mathbb{C}}^n$ of dimension ≥ 1 , every meromorphic function ξ defined in a complex connected neighborhood U of Y in $\mathbb{P}_{\mathbb{C}}^n$ can be (uniquely) extended to a meromorphic (and hence, rational) function on $\mathbb{P}_{\mathbb{C}}^n$. Severi proved this result in [Sev32] in the case when Y is a nonsingular hypersurface of $\mathbb{P}_{\mathbb{C}}^n$, and Barth generalised it to every closed connected subset Y of $\mathbb{P}_{\mathbb{C}}^n$ of positive dimension in [Bar68].

As we shall see in Proposition 1, the pairs (X, Y) as in the statement of Theorem 1 share a fundamental property. They are *universally connected*, that is:

Definition 2. Let X be a variety over an algebraically closed field k , and let Y be a closed subvariety of X . We say that the pair (X, Y) is *universally connected* if $f^{-1}(Y)$ is connected in X' for every proper surjective morphism $f: X' \rightarrow X$ from an irreducible variety X' .

According to a result of Bădescu and Schneider (see [BS02, (2.7)], cf. also [Băd04, 9.22]), universally connected pairs can be characterized by means of formal rational functions as follows:

Theorem 4. *Let X be an irreducible variety, and let Y be a closed subvariety of X . the following conditions are equivalent:*

- (i) (X, Y) is universally connected,
- (ii) $K(X/Y)$ is a field and $K(X)$ is algebraically closed in $K(X/Y)$,
- (iii) $K(X/Y)$ is a field and the algebraic closure of $K(X)$ in $K(X/Y)$ is purely inseparable over $K(X)$.

Remark 3. By Example 1, b) we see that if (X, Y) is $G3$ (with X irreducible) then $(X', f^{-1}(Y))$ is still $G3$, hence by Remark 1, a), (X, Y) is universally connected. Notice also that, If (X, Y) is a universally connected pair, then Y is $G3$ in X if and only if Y is $G2$ in X .

Examples 2. Let us recall some basic known examples and results.

- a) A point $Y = \{P\}$ in $\mathbb{P}^n$ is never $G2$ (and in particular, nor $G3$) by Example 1, a).
- b) Let $f: X' \rightarrow X$ be a proper surjective morphism of irreducible algebraic varieties, and let $Y \subseteq X$ and $Y' \subseteq X'$ be closed subvarieties such that $f(Y') \subseteq Y$. Assume that $K(X/Y), K(X'/Y')$, and $K(X'_{/f^{-1}(Y)})$ are fields. If Y' is $G3$ in X' , then Y is $G3$ in X ([Băd04, 9.23]).
- c) If Y' is $G2$ in X' , and if $f: X' \rightarrow X$ is a non constant dominant morphism of irreducible algebraic varieties, then $\dim f(Y') > 0$.
- d) Bădescu (see [Băd96, (0.1)], cf. also [Băd04, 11.1]) proved the following strengthening (and generalization) of Fulton-Hansen connectedness Theorem ([FH79]): for any proper morphism $f: X' \rightarrow \mathbb{P}^n(e) \times \mathbb{P}^n(e)$ from an irreducible variety X' , such that $\dim f(X) > n$, then $f^{-1}(\Delta_{\mathbb{P}^n(e)})$ is $G3$ in X' . Here $\mathbb{P}^n(e)$ denotes the n -dimensional weighted projective space of weights $e = (e_0, \dots, e_n)$, with $e_i \geq 0$, $i = 0, \dots, n$. Theorem 1 above plays an important role in the proof of this result.

Both our proofs follow the pattern suggested by the two facts below. The first shows, as promised, that the pairs (X, Y) as in Theorem 1 are universally connected.

Proposition 1. *Let Y be a closed subvariety of a projective irreducible subvariety X . Assume that $X \subset \mathbb{P}^n$, $\dim X = d > 2$ and Y is a set-theoretic intersection of X with r hyperplanes of $\mathbb{P}^n$, with $r \leq d - 1$. Then (X, Y) is universally connected.*

Proof. Let $f: X' \rightarrow X$ be a proper surjective morphism. We have to prove that $f^{-1}(Y)$ is connected. By the Stein factorization we may assume that f is finite. In this case $f^*(\mathcal{O}_X(1))$ is ample (and generated by its global sections) in which case the proposition follows from a result of Grothendieck (see [Gro05, Exposé XIII, Corollaire 2.2] (cf. also [Băd04, Ch. 7, Corollary 7.7])). $\square$

A way to prove that a universally connected pair (X, Y) is $G3$ is suggested by the following simple observation (cf. [Băd04, Proposition 9.23]):

Lemma 1. *Let X be an irreducible projective variety, and let Y be a closed subvariety of X . Assume (X, Y) is universally connected. Then (X, Y) is $G3$ if and only if there is a surjective proper morphism $f: X' \rightarrow X$ and a closed subvariety $Y'' \subset X'$ such that $f(Y'') \subset Y$ and (X', Y'') is $G3$.*

Proof. The necessity is trivial. For the converse: being (X, Y) universally connected $K(X'_{/f^{-1}(Y)})$ is a field, hence, using part b) of Remarks 1 we get that $(X', f^{-1}(Y))$ is $G3$. Then, by Example 2, b), (X, Y) is $G3$. $\square$

2. FIRST PROOF OF THEOREM 1

We show that an idea of Hironaka and Matsumura to prove the theorem in the case when $\dim(Y) = \dim(X) - r$ (i.e. if Y is a complete intersection in X , see [HM68, (4.3)]) can be suitably modified to yield a proof of Theorem 1 in general. Our surjective proper morphism, as in Lemma 1, is going to be a projection to X from the closure of the graph of a suitable linear projection of X .

Let $H_1, \dots, H_r$ in $\mathbb{P}^n$ be the hyperplanes cutting Y on X in $\mathbb{P}^n$, i.e. $Y = X \cap H_1 \cap \dots \cap H_r$ (set-theoretically); we can always assume that the H_i are all distinct.

Let us consider the $(n - r)$ -plane $H = H_1 \cap \dots \cap H_r \supseteq Y$ and let $L \subset H$ be a $(n - d - 1)$ -plane chosen in such a way that L does not contain any irreducible component of Y . Choose a d -plane $M \cong \mathbb{P}^d$, disjoint from L , and let us consider the linear projection:

$$\pi: \mathbb{P}^n \dashrightarrow M,$$

of center L . Setting $U := \mathbb{P}^n \setminus L$ we find a morphism $\pi_U: U \rightarrow M$. Note that $X_U := X \cap U \neq \emptyset$ is an open (and so dense) subset of X , and let us consider the morphism:

$$g_U := (\pi_U)|_{X_U}: X_U \rightarrow M,$$

together with its graph:

$$\Gamma_U = \{(x, y) \in X_U \times M \mid y = g_U(x)\} \subseteq X_U \times M \subseteq X \times M.$$

Let Γ be the closure of Γ_U in $X \times M$. So we get the commutative diagrams:

$$\begin{array}{ccc} \Gamma_U & \xrightarrow[(p_1)_U]{\cong} & X_U \\ (p_2)_U \downarrow & \swarrow g_U & \\ M & & \end{array} \qquad \begin{array}{ccc} X' := \Gamma & \xrightarrow{f:=p_1} & X \\ p_2 \downarrow & \searrow g & \\ M & & \end{array}$$

in which $(p_1)_U$ is an isomorphism and $(p_2)_U$ is dominant by the choice of L . So, since p_1 and p_2 are projective morphism, we see that p_1 and p_2 are surjective. Moreover, being $\Gamma_U \cong X_U$, we have that Γ is irreducible and p_1 is a birational regular map, this is our morphism f as in Lemma 1, and $X' = \Gamma$.

Now we need to find a closed subvariety $Y'' \subset X' = \Gamma$ which is G3 in X' such that $f(Y'') \subset Y$. To this end, let us consider $Y' := \pi_U(H \cap U) = H \cap M$ (because $H \supset L$). It is a $d-r$ -plane in $M \cong \mathbb{P}^d$, and since $d-r \geq 1$, we infer that Y' is G3 in M by Theorem 3. Therefore $Y'' := p_2^{-1}(Y')$ is G3 in Γ by Theorem 2. Clearly, by construction: $f(Y'') = p_1(p_2^{-1}(H \cap M)) \subseteq H \cap X = Y$.

Now, since Y'' is G3 in X' , by Lemma 1 and Proposition 1, Y is G3 in X . This finishes the first proof of Theorem 1.

3. SECOND PROOF OF THEOREM 1

This second proof makes use of a suitable incidence variety. Specifically, under the notation of Theorem 1, let $h_1, \dots, h_r$ in $\mathbb{P}^n$ be linear forms defining the distinct hyperplanes $H_1, \dots, H_r$ such that, set-theoretically $Y = X \cap H_1 \cap \dots \cap H_r$. Recall that by hypothesis we have $r+1 \leq d = \dim(X)$. Let us consider the projective space:

$$P := \mathbb{P}(H^0(\mathcal{O}_{\mathbb{P}^n}^{\oplus(r+1)}(1))) \cong \mathbb{P}^{(n+1)(r+1)-1}.$$

Taken a global section $\sigma \in H^0(\mathcal{O}_{\mathbb{P}^n}(1))$, let us denote by $V(\sigma)$ the zero locus of σ in $\mathbb{P}^n$, which is a hyperplane in $\mathbb{P}^n$ in the case that σ

is nonzero, and coincides with $\mathbb{P}^n$ otherwise. Let $Z \subset X \times P$ be the closed incidence subvariety given by:

$$Z \{ (x, [\sigma_0, \dots, \sigma_r]) \in X \times P \mid x \in X \cap V(\sigma_0) \cap \dots \cap V(\sigma_r) \}.$$

Let us consider the two projections:

$$\begin{array}{ccc} Z & \xrightarrow{f} & X \\ g \downarrow & & \\ P & & \end{array}$$

and note that f is surjective and all its fibres are linear subspaces of P of the same dimension. Since X is irreducible, we deduce that Z is also irreducible. The proper morphism g is also surjective because $r + 1 \leq d$, and hence $X \cap V(\sigma_0) \cap \dots \cap V(\sigma_r) \neq \emptyset$ for each point $[\sigma_0, \dots, \sigma_r] \in H^0(\mathcal{O}_{\mathbb{P}^n}^{\oplus(r+1)}(1)) \setminus \{0\}$. Moreover,

$$g^{-1}([\sigma_0, \dots, \sigma_r]) = Y \times \{[\sigma_0, \dots, \sigma_r]\} \subset f^{-1}(Y)$$

for each point $[\sigma_0, \dots, \sigma_r] \in H^0(\mathcal{O}_{\mathbb{P}^n}^{\oplus(r+1)}(1))$ such that $\sigma_0, \dots, \sigma_r$ generate the same vector space as $h_1, \dots, h_r$.

As in the first proof, the theorem will be proved once we show that $g^{-1}(L) \subseteq f^{-1}(Y)$ for some connected closed subset $L \subset P$, with $\dim(L) \geq 1$. We show that we can take for L a suitable line. Let us consider the points of P defined by:

$$q_1 := [h_1, \dots, h_r, 0] \text{ and } q_2 := [0, h_1, \dots, h_r],$$

and let us denote by $L \subset P$ the line joining the two points. Since any point of L is of type $\lambda q_1 + \mu q_2 = [\lambda s_1, \lambda s_2 + \mu s_1, \dots, \lambda s_r + \mu s_{r-1}, \mu s_r]$, we easily see that:

$$g^{-1}(L) = Y \times L \subseteq f^{-1}(Y) \subset Z.$$

This concludes the second proof.

REFERENCES

- [Băd96] Lucian Bădescu, *Algebraic Barth-Lefschetz theorems*, Nagoya Math. J. **142** (1996), 17–38.
- [Băd04] ———, *Projective geometry and formal geometry*, Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) [Mathematics Institute of the Polish Academy of Sciences. Mathematical Monographs (New Series)], vol. 65, Birkhäuser Verlag, Basel, 2004.
- [Bar68] Wolf Barth, *Fortsetzung, meromorpher Funktionen in Tori und komplex-projektiven Räumen*, Invent. Math. **5** (1968), 42–62.

- [BS02] Lucian Bădescu and Michael Schneider, *Formal functions, connectivity and homogeneous spaces*, Algebraic geometry, de Gruyter, Berlin, 2002, pp. 1–23.
- [Fal80] Gerd Faltings, *A contribution to the theory of formal meromorphic functions*, Nagoya Math. J. **77** (1980), 99–106.
- [FH79] William Fulton and Johan Hansen, *A connectedness theorem for projective varieties, with applications to intersections and singularities of mappings*, Ann. of Math. (2) **110** (1979), no. 1, 159–166. MR MR541334 (82i:14010)
- [Gro60] A. Grothendieck, *Éléments de géométrie algébrique. I. Le langage des schémas*, Inst. Hautes Études Sci. Publ. Math. (1960), no. 4, 228.
- [Gro61] ———, *Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I*, Inst. Hautes Études Sci. Publ. Math. (1961), no. 11, 167.
- [Gro05] Alexander Grothendieck, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 4, Société Mathématique de France, Paris, 2005, Séminaire de Géométrie Algébrique du Bois Marie, 1962, Augmenté d’un exposé de Michèle Raynaud. [With an exposé by Michèle Raynaud], With a preface and edited by Yves Laszlo, Revised reprint of the 1968 French original.
- [Har70] Robin Hartshorne, *Ample subvarieties of algebraic varieties*, Notes written in collaboration with C. Musili. Lecture Notes in Mathematics, Vol. 156, Springer-Verlag, Berlin, 1970.
- [Har77] ———, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.
- [HM68] Heisuke Hironaka and Hideyuki Matsumura, *Formal functions and formal embeddings*, J. Math. Soc. Japan **20** (1968), 52–82.
- [Sev32] F. Severi, *Alcune proprietà fondamentali dell’insieme dei punti singolari di una funzione analitica di più variabili.*, Memorie Reale Acad. d’Italia, Roma, 3, Mat. Nr. 1, 20 p , 1932 (Italian).

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DEGLI STUDI
DI CATANIA, VIALE A. DORIA 6, 95125 CATANIA
E-mail address: bonacini@dmf.unict.it

DIPARTIMENTO DI MATEMATICA, VIA DODECANESO 35, 16146 GENOVA
E-mail address: delpadro@dima.unige.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI ROMA TRE, LARGO
SAN LEONARDO MURIALDO 1, 00146 ROMA
E-mail address: nesci@mat.uniroma3.it